arXiv:hep-th/9302117v1 24 Feb 1993

UCI-TR 93-6

Equations of Motion for Spinning Particles in External Electromagnetic and Gravitational Fields

Karl Yee and Myron Bander*

Department of Physics, University of California, Irvine, California 92717, USA

(Received February 1993)

The equations of motion for the position and spin of a classical particle coupled to an external electromagnetic and gravitational potential are derived from an action principle. The constraints insuring a correct number of independent spin components are automatically satisfied. In general the spin is not Fermi-Walker transported nor does the position follow a geodesic, although the deviations are small for most situations.

PACS Numbers: 03.20.+i, 04.20.Fy

Typeset Using *REVTEX*

The derivation of the equations of motion of *classical* spinning particles in external fixed fields has occupied physicists for over fifty years. In special relativity it was first attacked by Frenkel [1]. Using his work as a basis Bargmann, Michel and Telegdi [2] discussed the precession of spinning particles in external electromagnetic fields. It is amusing to note that, even today, there is a controversy as to the torque and force on such particles in space and time dependent fields [3]. The discussion of a spinning particle in an external gravitational potential goes back to Papapetrou [4] who endowed a particle with spin by considering a rotating mass-energy distribution in the limit of vanishing volume but with the angular momentum remaining finite. Results were obtained using Grassmann variables and supersymmetry [5] and an attempt at a general procedure was made by Khriplovich [6]. Most of the emphasis in the above works is on the equations for the spin components and, with the exception of Refs. [1,3,4], the equations for the motion of the position of the particle are either ignored or are incomplete. We shall present a canonical procedure for obtaining the equations of motion, both, for the spin and position. The results are identical to those that would have been obtained using the method of Ref. [1].

We shall obtain the equations of motion in terms of the proper time, τ, of the particle. The position of the particle will be denoted by x^μ and the spin will be described by the antisymmetric spin matrix S_{ab}. As usual, Greek indices will denote covariant vectors, tensors, etc. and Latin ones those in local Lorentz frames; these are connected by the vierbein field, $e^a_\mu(x)$. The spin matrix satisfies the Poisson relation

$$\{S_{ab}, S_{cd}\} = \eta_{ac} S_{bd} + \eta_{bd} S_{ac} - \eta_{ad} S_{bc} - \eta_{bc} S_{ad}\,, \tag{1}$$

with η_{ab} the flat space metric. It will prove to be convenient to obtain the equations of motion for the position from a Lagrangian and those for the spin from a Hamiltonian; such a combined procedure calls for the introduction of a Routhian [7] $\mathcal{R}(x^\mu, S_{ab})$. At the end of this work we shall provide an expression for the action. The equations of motion are

$$\begin{aligned} \frac{\delta \int d\tau \mathcal{R}}{\delta x^\mu} &= 0\,, \\ \frac{dS_{ab}}{d\tau} &= \{\mathcal{R}, S_{ab}\}\ ; \end{aligned} \tag{2}$$

τ is the proper time. All the variables we have considered are not independent, but satisfy various constraints. With u^μ, the four velocity, these are

$$u^\mu u_\mu = 1\,, \tag{3}$$

$$S_{ab}S^{ab} = 2s^2\,, \tag{4}$$

$$S_{ab}u^a = 0\,. \tag{5}$$

Eq. (3) guarantees that τ is the proper time and will be satisfied as long as $\mathcal{R}$ is written in a reparametrization invariant form and Eq. (5) is satisfied. Eq. (4) insures that the spin of the particle is constant and is equal to s; it is automatically satisfied for all situations considered. Eq. (5) results from the fact that in the particle's rest frame the spin tensor has only three independent components; it is this constraint that causes all the complications.

For a particle in an external field derived from a vector potential A_μ and a gravitational field specified by the spin connection ω_μ^{ab} a tempting Routhian is

$$\mathcal{R}_0 = m\sqrt{u^2} + eu^\mu A_\mu - \frac{u^\mu}{2}\omega_\mu^{ab}S_{ab} + \frac{eg}{4m}F^{ab}S_{ab}\sqrt{u^2} - \frac{\kappa}{8m}R^{abcd}S_{ab}S_{cd}\sqrt{u^2}\,. \tag{6}$$

m is the mass of the particle and e is its charge; g is the gyromagnetic ratio and κ specifies the strength of gravitational magnetic moment coupling introduced in Ref. [6]. This is the most general Routhian not involving derivatives of the field strength tensor or of the Riemann tensor and not involving terms of the form $S_{ab}u^a$. Unfortunately, the equations of motion derived using $\mathcal{R}_0$ do not satisfy the constraints of Eq. (5). There are two procedures that will guarantee this constraint; both give the same equations of motion. One can follow the method of Ref. [1] and add to $\mathcal{R}_0$ the constraint multiplied by a Lagrange multiplier. We shall follow a different procedure. First define

$$\tilde{S}_{ab} = S_{ab} - S_{ac}\frac{u^c u_b}{u^2} - S_{cb}\frac{u^c u_a}{u^2}\,. \tag{7}$$

The Poisson brackets of the $\tilde{S}_{ab}$'s is the one given in Eq. (1) with the metric tensor η_{ab} replaced by $\eta_{ab} - u_a u_b/u^2$. The $\tilde{S}_{ab}$ are related to the spin vector s^a by

$$
\begin{aligned}
s^a &= -\frac{1}{2}\epsilon^{abcd}\tilde{S}_{bc}u_d\,, \\
\tilde{S}_{ab} &= \epsilon_{abcd}s^c u^d\,.
\end{aligned} \tag{8}
$$

The spin vector satisfies $s^a s_a = s^2$ and the constraint $s^a u_a = 0$.

A desired Routhian is obtained by replacing all S_{ab}'s in Eq. (6) by $\tilde{S}_{ab}$'s and by adding $(du^a/d\tau)\, S_{ab}u^b/u^2$ to it;

$$
\begin{aligned}
\mathcal{R} = m\sqrt{u^2} + eu^\mu A_\mu - \frac{u^\mu}{2}\omega_\mu^{ab}S_{ab} + \frac{Du^a}{D\tau}\frac{S_{ab}u^b}{u^2} \\
+ \frac{eg}{4m}F^{ab}\tilde{S}_{ab}\sqrt{u^2} - \frac{\kappa}{8m}R^{abcd}\tilde{S}_{ab}\tilde{S}_{cd}\sqrt{u^2}\,.
\end{aligned} \tag{9}
$$

Here, D denotes a covatiant derivative and we have used the identity

$$
\frac{du^a}{d\tau}\frac{S_{ab}u^b}{u^2} - \frac{u^\mu}{2}\omega_\mu^{ab}\tilde{S}_{ab} = \frac{Du^a}{D\tau}\frac{S_{ab}u^b}{u^2} - \frac{u^\mu}{2}\omega_\mu^{ab}S_{ab}\,. \tag{10}
$$

The fourth term in Eq. (9) involves the acceleration explicitly; it adds the Thomas precession term to the equations of motion for the spin and insures that $S_{ab}u^a$ may be set equal to zero.

The equations of motion, with Eqs. (3-5) satisfied, for the spin tensor are

$$
\begin{aligned}
\frac{DS_{ab}}{D\tau} + (u_b S_{ac} - u_a S_{bc})\,\frac{Du^c}{D\tau} = -\left(\frac{eg}{2m}F^{cd} + \frac{\kappa}{2m}R^{cdef}S_{ef}\right) \\
\times\left[S_{ac}(\eta_{bd} - u_b u_d) + S_{bd}(\eta_{ac} - u_a u_c)\right]\,.
\end{aligned} \tag{11}
$$

This expression is consistent with Eq. (4) and Eq. (5). The equations of motion for the coordinates of the particle are

$$
\begin{aligned}
m\tfrac{Du_\mu}{D\tau} - eF_{\mu\nu}u^\nu - \tfrac{1}{2}R^{cd}_{\mu\nu}S_{cd}u^\nu = \tfrac{D}{D\tau}\left[\left(-\tfrac{eg}{4m}F^{cd}S_{cd} + \tfrac{\kappa}{8m}R^{cdef}S_{cd}S_{ef}\right)u_\mu\right] \\
\tfrac{D}{D\tau}\left(e^c_\mu S_{cd}\tfrac{Du^d}{D\tau}\right) + \tfrac{eg}{4m}F^{cd}{}_{;\mu}S_{cd} - \tfrac{\kappa}{8m}R^{cdef}{}_{;\mu}S_{cd}S_{ef} \\
+ \tfrac{D}{D\tau}\left[e^h_\mu\left(\tfrac{eg}{2m}F^{cd}u_c S_{hd} - \tfrac{\kappa}{2m}R^{cdef}S_{cd}u_e S_{hf}\right)\right]\,.
\end{aligned} \tag{12}
$$

These equations are exact. Except in the case of large gravitational field gradients the modifications due to the spin will be small [8]. It is interesting to study various limits of Eq. (12). If we ignore the right hand side of that equation and plug the results into Eq. (11) we obtain

$$\begin{aligned}\frac{DS_{ab}}{D\tau} = &-\left(\frac{eg}{2m}F^{cd} + \frac{\kappa}{2m}R^{cdef}S_{ef}\right)(S_{ac}\eta_{bd} + S_{bd}\eta_{ac}) \\ &+\left[\frac{e(g-2)}{2m}F^{cd} + \frac{(\kappa-1)}{2m}R^{cdef}S_{ef}\right](S_{ac}u_b u_d + S_{bd}u_a u_c)\ . \end{aligned} \tag{13}$$

We know that the electromagnetic part of the equations of motion for the spin simplify in the case $g = 2$; we also see that there is a simplification for the gravitational part in the case $\kappa = 1$. That the Dirac equation

$$\gamma^a e^\mu_a\left(i\partial_\mu - eA_\mu + \frac{1}{2}\omega^{cd}_\mu S_{cd}\right)\psi - m\psi = 0\,, \tag{14}$$

with S_{cd} expressed in terms of the Dirac γ matrices, yields $g = 2$ is well known; it also yields $\kappa = 1$. We note that even in the presence of only gravitational couplings, but with $\kappa \neq 0$, the spin is *not* Fermi-Walker transported [8,9]. The corrections to Fermi-Walker transport are very small, except as mentioned earlier, in the presence of large gravitational fields and gradients.

Another interesting limit is the situation of no electromagnetic field and $\kappa = 0$. The equation of motion for the position of the particle is

$$m\frac{Du_\mu}{D\tau} - \frac{1}{2}R^{cd}_{\mu\nu}S_{cd}u^\nu - \left(e^c_\mu S_{cd}\frac{D^2u^d}{D\tau^2}\right) = 0\,. \tag{15}$$

This agrees with the equations in Ref. [4]; we note that spinning particles do not follow geodesics. The term involving the time derivative of the acceleration has been interpreted, in Ref. [4], as being responsible for classical *Zitterbewegung*; in the following sense we agree with this interpretation: in the absence of gravitational interactions a spinning particle will oscillate in the plane perpendicular to the spin direction with a frequency $\omega = E/s$; $s^2 = S^{ab}S_{ab}/2$ is the magnitude of the spin vector. If we set $|s| = \hbar/2$ we recover the quantum mechanical *Zitterbewegung* frequency.

In the non-relativistic limit Eq. (12), for a purely magnetic dipole interaction is

$$m\frac{d\boldsymbol{v}}{dt} = \frac{eg}{2m}\boldsymbol{\nabla}(\boldsymbol{H}\cdot\boldsymbol{s}) + \frac{eg}{2m}\frac{d}{dt}(\boldsymbol{E}\times\boldsymbol{s})\ ; \tag{16}$$

s is defined in Eq. (8). This expression agrees with the force equation advocated in Ref. [3].

For completeness we present an action which corresponds to the Routhian of Eq. (9). A convenient approach is to add a Wess-Zumino [10] term. For closed paths in proper time we introduce a two dimensional manifold, $\mathcal{M}$, parametrized by $y_1,\ y_2$ whose boundary is the path τ. The action is

$$\mathcal{A} = \frac{1}{s^2} \int_{\mathcal{M}} dy_1\ dy_2 \epsilon^{\alpha\beta} \mathrm{Tr} S \frac{\partial S}{\partial y_\alpha} \frac{\partial S}{\partial y_\beta} + \int d\tau \mathcal{R} \,. \tag{17}$$

The canonical equations obtained from the double integral yield the Poisson brackets of Eq. (1)

We wish to thank Professor A. Schwimmer for discussions on the Wess-Zumino term. This work was supported in part by the National Science Foundation under Grant No. PHY-9208386.

REFERENCES

* e-mail: mbander@funth.ps.uci.edu

[1] J. Frenkel, Z. Physik **37**, 243 (1926).

[2] V. Bargmann, L. Michel and V. L. Telegdi, Phys. Rev. Lett. **2**, 435 (1959).

[3] Y. Aharonov and A. Casher, Phys. Rev. Lett. **53**, 319-321 (1984); L. Vaidman, Am. J. Phys. **58**, 978 (1986).

[4] A. Papapetrou, Proc. Roy. Soc. **A209**, 248 (1951).

[5] A. Barducci, R. Casalbuoni and L. Lusanna, Nuovo Cimento **35A**, 389 (1976); F. Ravndal, Phys. Rev. D**21**, 2823 (1980).

[6] I. B. Khriplovich, Zh. Exsp. Teor. Fiz. **96**, 385 (1989) [Sov. Phys. JETP **69** 217 (1989)].

[7] L. D. Landau and E. M. Lifshitz, *Classical Mechanics* (Pergamon Press, Oxford, 1960) p. 134.

[8] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation* (W. H. Freeman and Company, San Francisco, 1973) p. 1121.

[9] S. Weinberg, *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (John Wiley and Sons, Inc. , New York, 1972), p. 121.

[10] J. Wess and B. Zumino, Phys. Lett. **37B**, 95 (1971).